

## Nutritional and Socioeconomic Status Toward The Low-Birth-Weight Neonates (LBW) Case

Hasriyani<sup>1</sup>, Suharyo Hadisaputro<sup>1</sup>, Karmila Budhi<sup>2</sup>, Mexitalia Setiawati<sup>2</sup>, Henry Setyawan<sup>1</sup>

<sup>1</sup>Department of Epidemiology, Postgraduate Program, Diponegoro University, Indonesia

<sup>2</sup>Dr. Kariadi General Hospital, Semarang, Indonesia

### Abstract

LBWN is a public health problem and is still a major cause of perinatal morbidity and mortality. In Indonesia, in the year of 2013, the case rate of LBWN is 10.2%. This study aims to prove the factor of nutritional status, socioeconomic status toward the case of LBWN. This research used case control study design. The total sample was 138 infants, consisting of 69 cases and 69 controls that meet the inclusion and exclusion criteria. Case collecting sample was conducted through consecutive sampling and simple random sampling control, by matching the sex of the baby and their area. Data analysis used logistic regression. The result of the analysis showed that the low of nutritional status (OR = 3.159; 95% CI = 1.390-7.178), low socioeconomic status (OR = 2.175; 95% CI = 1.066-4.439) were risk factors for LBWN case. Low nutritional status and low socioeconomic status are risk factors for LBWN case. It is needed to provide CIE (Counseling, Information, and Education) intensively to pregnant women and expectant pregnant women related to good nutrition.

**Keywords:** LBWN, risk factor, nutritional status, socioeconomic status.

### I. INTRODUCTION

Low-birth-weight Neonates (LBWN) is defined as a newborn which is born with a birth weight of <2500 grams.<sup>(1,2)</sup> LBWN is still a health problem in many countries because it is considered to be one of the factors causing infant mortality.<sup>(3)</sup> LBWN case according to WHO is 15.5% of 20 million live births per year, 96.5% is in developing countries and contributes 60%-80% of all neonatal deaths.<sup>(2,4,5)</sup>

Early neonatal mortality (age 0-6 days) in Indonesia in 2007 was 78.5% with the cause of death was LBWN (32.4%). Infant Mortality Rate (IMR) of 32 deaths per 1000 live births. Neonatal Mortality Rate (NMR) of 19 per 1000 live births (the same number as in 2007). The attention of efforts to decrease the NMR (0-28 days) becomes important because neonatal mortality contributes to 56% of infant mortality.<sup>(6)</sup> LBWN case in 2013 is 10.2%. This number decreased if it's compared to the 2010 of 11.1%.<sup>(7)</sup>

Nationally, South Sulawesi Province is one of the provinces with the 7th highest percentage of LBWN (12.4%).<sup>(8)</sup> In 2014, Makassar is one of the highest LBW case among 24 regencies/cities in South Sulawesi (2.48%).<sup>(9)</sup>

The proportion of LBWN cases in Makassar City in 2014 was 2.80% with 2.46% mortality, 2.62% in 2015 with 1.96% mortality and 3.29% of LBWN cases in 2016. From 46 Public Health Center (PHC) in Makassar City, there were 6 health centers with highest percentage of LBWN is 12.19% of Antang Public Health Center, Tamangapa Public Health Center is 10.47%, Kaluku Bodoa Public Health Center 9.46%, Minasa Upa Public Health Center 8.64%, Rappokalling Public Health Center 7.44%, Patingalloang Public Health Center 7.24%.<sup>(10)</sup>

LBWN is the result of the interaction of various risk factors such as maternal factors (maternal age, pregnancy range, extent, pregnancy diseases, maternal education, socioeconomic status, nutritional status, antenatal care status, nutritional intake), placental factors, fetal factors, and environmental factors.<sup>(11-15)</sup>

Based on the high rate of LBWN in Makassar, it is necessary to see the risk factors that can cause the occurrence of LBWN. The purpose of this study is to prove the factor of nutritional status and socioeconomic status toward the LBWN case.

### II. METHOD

The type of research was observational analytic research with case control design. Population of group case study in this research is all babies born with <2500 gram birth weight at Antang Public Health Center, Tamangapa Public Health Center, Kaluku Bodoa Public Health Center, Minasa Upa Public Health Center, Rappokaling Public Health Center and Patingalloang Public Health Center of Makassar City registered in register book of cohort mother and/or register book of postpartum in year 2016/2017. While the control group study population were all babies born with birth weight  $\geq$ 2500 gram at Antang Public Health Center, Tamangapa Public Health Center, Kaluku Bodoa Public Health Center, Minasa Upa Public Health Center, Rappokaling Public Health Center, Patingalloang Public Health Center of Makassar City registered in register book of cohort mother and / or register book of postpartum in year 2016/2017. Collecting sample of cases was done by of consecutive sampling and was controlled by simple random sampling by matching the sex of the baby and the area. Dependent variable in this

research is LBWN. The independent variables were maternal age, nutritional status, pregnancy range, Antenatal Care (ANC) and socioeconomic status. Confounding variables were maternal education levels and parity. Data analysis conducted in this research was univariate, bivariate and multivariate analysis at significance level  $p = <0.05$ . This research has got the research ethic approval from Health Research Ethics Commission (KEPK) Faculty of Medicine, Diponegoro University and dr. Kariadi Hospital Semarang with No. 361 / EC / FK-RSDK / VI / 2017.

### III. RESULT

Distribution of maternal age in case group was mostly with age of 24-27 years old, the youngest age was 15 years old to the oldest 43 years old with average age of 26 years. While the control group was mostly in the age group of 24-27 years old, the youngest age of 16 years old to the oldest 45 years old with an average age of 27 years.

Table 1. Distribution of Respondent Characteristics based on Maternal Age, Educational Background, Occupation, Pregnancy Age and Newborn's Sex

Respondent Characteristics	LBWN status			
	Case		Control	
	n	(%)	n	(%)
1. Maternal age when gave birth				
12-15 years old	1	1.4	0	0
16-19 years old	9	13.1	9	13.1
20-23 years old	13	18.8	12	17.4
24-27 years old	21	30.4	16	23.1
28-31 years old	15	21.7	9	13.1
32-35 years old	4	5.8	9	13.1
36-39 years old	3	4.4	8	11.6
40-43 years old	3	4.4	5	7.2
44-47 years old	0	0	1	1.4
Total	69	100.0	69	100.0
Mean	26.07		27.9	
Median	26.0		27.0	
Modus	27		21	
Minimum	15		16	
Maximum	43		44	
2. Educational Background				
None	0	0.0	1	1.4
Elementary	20	29.0	24	34.8
Junior High School	17	24.6	20	29.1
Senior High School	24	34.8	18	26.1
Diploma	1	1.4	3	4.3
Bachelor	7	10.1	3	4.3
Total	69	100.0	69	100.0
3. Occupation				
Housewife	57	82.6	64	92.8
Employee	0	0.0	0	0.0
Farmer	0	0.0	0	0.0
Fisherman	0	0.0	0	0.0
Civil Employee	0	0.0	0	0.0
Private Employee	6	8.7	2	2.9
Entrepreneur	1	1.4	3	4.3
Other	5	7.2	0	0.0
Total	69	100.0	69	100.0
4. Pregnancy Age				
Premature (<37 weeks)	14	20.3	2	2.9
Exact month (37-41 weeks)	55	79.7	67	97.1
Total	69	100.0	69	100.0
5. Newborn Sex				
Male	35	50.7	35	50.7
Female	34	49.3	34	49.3
Total	69	100.0	69	100.0

Table 2 shows the independent variables or the main variables that have significant relationship with the LBWN case ( $p < 0.05$ ) that was nutritional status ( $p = 0.004$ ; OR = 3.188; 95% CI = 1.421-7.151); and socioeconomic status ( $p = 0.025$ ; OR = 2.199; 95% CI = 1.101-4394).

Table 2. Bivariate Analysis of Independent and Confounding Variables

No	Variable	Case		Control		OR	95% CI	p
		n	%	n	%			
1.	Maternal age							
	- Risky	16	23.2	23	33.3	0.604	0.285-1.279	0.186
	- Not risky	53	76.8	46	66.7			
2.	Nutritional status							
	- Low	58	84.1	43	62.3	3.188	1.421-7.151	0.004
	- High	11	15.9	26	37.7			
3.	Pregnancy Range							
	- Risky	5	7.2	9	13.0	0.521	0.165-1.643	0.259
	- Not risky	64	92.8	60	87.0			
4.	ANC status							
	- Incomplete	38	55.1	38	55.1	1.000	0.511-1.956	1.000
	- Complete	31	44.9	31	44.9			
5.	Socioeconomic status							
	- Low	47	68.1	34	49.3	2.199	1.101-4.394	0.025
	- High	22	31.9	35	50.7			
6.	Mother's education							
	- Low	37	53.6	45	65.2	0.617	0.311-1.223	0.165
	- High	32	46.4	24	34.8			
7.	Parity							
	- Risky	38	55.1	38	55.1	1.000	0.511-1.956	1.000
	- Not risky	31	44.9	31	44.9			

Table 3 shows the independent factors that were risk factors for LBWN case in the multivariate model was low nutritional status and low socioeconomic status.

Table 3. Result of Logistic Regression Analysis

No	Risk factors	B value	OR	95%CI	P
1.	Low nutritional status	1.150	3.159	1.390-7.178	0.006
2.	Low socioeconomic status	0.777	2.175	1.066-4.438	0.033
	Constanta	-1.311			

## IV. DISCUSSION

### A. Nutritional Status

The nutritional status of pregnant women as measured by weight gain during pregnancy and Measuring Upper Arm Circle (MUAC) is positively related to the energy consumption level of pregnant women. Low of nutrition will cause bad consequences for the mother and her fetal and can cause fetal growth inhibited and LBWN.<sup>(16,17)</sup> Based on interviews with mothers, it was found that 52.2% of bad MUAC and weight gain were 79.7%. Multivariate analysis showed that low nutritional status was a risk factor for LBWN case with OR = 3.159 (95% CI = 1,390-7,178), meaning that low nutritional status was 3.159 times higher for LBWN delivery compared with good nutritional status. This result is consistent with the research of Tazkiah et al (2013) indicated that mother's nutritional status is a risk factor to the case of LBWN (OR = 2.583; 95% CI = 1.269-2.257).<sup>(16)</sup> Research from Reza (2014) showed that MUAC <23.5 cm have risk to the LBWN case (OR=3.678; CI95%=1.125-12.027).<sup>(18)</sup> Mumbare et al (2012) study showed that body weight during pregnancy was a risk factor for the case of LBW (OR = 4.82; CI95% = 2.54-9.15 ).<sup>(19)</sup>

The determinants of LBWN are not only during pregnancy, but also before pregnancy, therefore the nutritional status of the mother before pregnancy should be in good condition. During pregnancy, the need for food substances increases so that when pregnancy nutritional intake must be adequate so as not to disturb the growth of the fetal. Some of the factors causing fetal growth disturbance related to the mechanism are maternal nutritional intake, nutritional supply to the uterus and placenta, transport of nutrition through the placenta, and regulation of fetal nutrition.<sup>(18)</sup>

### B. Socioeconomic Status

Income has an indirect effect on the case of LBWN. Families with high income will be able to meet the nutritional needs, otherwise families with low income will have difficulties in meeting the nutritional needs.<sup>(3)</sup> Low economic level is one of the factors that cause the decline in affordability of food to meet the needs that affect the quality and quantity of food consumed by the whole family. If it continues, the nutritional status of the family, especially pregnant women, will be worse.<sup>(20)</sup>

Multivariate analysis showed that low socioeconomic status was a risk factor for LBWN case with OR = 2,175 (95% CI = 1,066-4,438), meaning low socioeconomic status had 2,213 times greater risk for delivering LBWN

than high socioeconomic status. In line with a research by Jayant et al, (2011) shows that socioeconomic status is a risk factor for LBWN case. (OR = 1.68; 95% CI = 1.04-2.71)<sup>(21)</sup>

## V. CONCLUSION

Based on the results of the study, it can be concluded that the low of nutritional status and low socioeconomic status are risk factors of the LBWN case.

### References

1. WHO. Global Nutrition Target 2025: Low Birth Weight Policy Brief. Geneva; 2014.
2. WHO. Optimal Feeding Of Low Birthweight Infants in Low-And Middle Income Countries. Geneva: World Health Organization; 2011. 1-51 p.
3. Pramono MS, Paramita A. Pattern of Occurrence and Determinants of Baby With Low Birth Weight in Indonesia 2013. *Bul Penelit Sist Kesehat*. 2015;18(1).
4. United Nations Children's Fund and World Health Organization. Low Birthweight: Country, regional and global estimates. Unicef. 2004. 1-31 p.
5. WHO. Maternal, Newborn, Child and Adolescent Health Care of the Preterm and/or Low-Birth-Weight Newborn. 2013.
6. Kemenkes RI. Propil Kesehatan Indonesia Tahun 2014. Jakarta: Kementerian Kesehatan Republik Indonesia; 2015. 106-107 p.
7. Kemenkes RI. Riset Kesehatan Dasar (RISKESDAS) Tahun 2013. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia; 2013. 182 p.
8. Kementerian Kesehatan RI. Riset Kesehatan Dasar Dalam Angka Provinsi Sulawesi Selatan Tahun 2013. Jakarta: Badan Penelitian dan Pengembangan Kesehatan; 2013. 1-425 p.
9. Dinkes Provinsi Sulawesi Selatan. Profil Kesehatan Sulawesi Selatan 2014. Makassar: Dinas Kesehatan Provinsi Sulawesi Selatan; 2015. 1-156 p.
10. Dinkes Kota Makassar. Profil Kesehatan Kota Makassar Tahun 2014. Makassar: Dinas Kesehatan Kota Makassar; 2015. 1-92 p.
11. Amiruddin R, Hasmi. Determinan Kesehatan Ibu dan Anak. 1st ed. Jakarta: CV. Trans Info Media; 2014. 1-217 p.
12. Mochtar AB. Persalinan Preterm. In: Rachimhadhi T, Wiknjastro GH, editors. Ilmu Kebidanan Sarwono Prawirohardjo. 4th ed. Jakarta: Bina Pustaka Sarwono Prawirohardjo; 2008. p. 667-79.
13. Manuaba AC, Manuaba IBG, Manuaba BG. Ilmu Kebidanan, Penyakit Kandungan, dan KB untuk Pendidikan Bidan. 2nd ed. Jakarta: EGC; 2010. 1-669 p.
14. Krisnadi SR. Faktor Risiko Persalinan Prematur. In: Krisnadi SR, Effendi JS, Pribadi A, editors. Prematritas. 1st ed. Bandung: PT. Refika Aditama; 2009. p. 43-65.
15. Gomella TL, Cunningham MD, Eyal FG. Neonatology Management, Procedures, On-Cell Problems, Diseases, and Drugs. Seventh Ed. America: MC Graw Hill Education Lange; 2013. 1-1059 p.
16. Tazkiah M, Wahyuni CU, Martini S. Epidemiological Determinants Low Birth Weight in Malaria Endemic Areas Banjar Distric. *J Berk Epidemiol*. 2013;1(2):266-76.
17. Yongky, Hardinsyah, Gulardi, Marham. Nutritional Status of First Trimester and the Accretion of Pregnant Women Body Weight and Its Relationship with Low Birth Weight. *J Gizi dan Pangan*. 2009;4(1):8-12.
18. Reza Chaerul, Puspitasari N. Determinants of Low-Birth-Weight-Neonates. *J Biometrika dan Kependud*. 2014;3(2):96-106.
19. Mumbare S, Maindarkar G, Darade R, Yenge S, Tolani MK, Patole K. Maternal Risk Factors Associated with Term Low Birth Weight Neonates : A Matched-Pair Case Control Study. *Indian Pediatr* Januari 16. 2012;49:25-8.
20. Ratnasari D, Suhartono, Rahfiludin MZ. Risk Factors for Low Birth Weight in the Agricultural Area (study in Brebes district). *J Gizi Pangan*. 2017;12(1):41-8.
21. Jayant D, Phalke, Bangal, Peeyuusha, Sushen B. Maternal Risk Factors for Low Birth Weight Neonates : a Hospital Based Case-Control Study in Rural Area of Western Maharashtra ., *Natl J Community Med*. 2011;2(3):394-8.